

CONTINUOUS-TRACE GROUPOID CROSSED PRODUCTS

IGOR FULMAN, PAUL S. MUHLY, AND DANA P. WILLIAMS

ABSTRACT. Let G be a second countable, locally compact groupoid with Haar system, and let $\mathcal{A}$ be a bundle of C^* -algebras defined over the unit space of G on which G acts continuously. We determine conditions under which the associated crossed product $C^*(G; \mathcal{A})$ is a continuous trace C^* -algebra.

1. INTRODUCTION

Throughout this note, G will denote a second countable, locally compact groupoid with Haar system $\{\lambda^u\}_{u \in G^{(0)}}$. Also, we shall fix a bundle¹ $\mathcal{A}$ of (separable) C^* -algebras over the unit space $G^{(0)}$ of G . We shall write p for the projection of $\mathcal{A}$ onto $G^{(0)}$. We shall assume that there is a continuous action, denoted σ , of G on $\mathcal{A}$. This means the following: First of all, σ must be a homomorphism from G into the isomorphism groupoid of $\mathcal{A}$, $\text{Iso}(\mathcal{A})$, so that, in particular, $\sigma_\gamma : A(s(\gamma)) \rightarrow A(r(\gamma))$ is a C^* -isomorphism for each $\gamma \in G$. Secondly, let $s^*(\mathcal{A})$ and $r^*(\mathcal{A})$ be the bundles on G obtained by pulling back $\mathcal{A}$ via s and r , so that $s^*(\mathcal{A}) = \{(\gamma, a) \mid a \in A(s(\gamma))\}$ and similarly for $r^*(\mathcal{A})$. Then σ determines a bundle map $\sigma^* : s^*(\mathcal{A}) \rightarrow r^*(\mathcal{A})$ by the formula $\sigma^*(\gamma, a) = (\gamma, \sigma_\gamma(a))$. The continuity assumption that we make is that for each continuous section f of $s^*(\mathcal{A})$, $\sigma^* \circ f$ is a continuous section of $r^*(\mathcal{A})$.

Let $C_c(G, r^*(\mathcal{A}))$ denote the space of continuous sections of $r^*(\mathcal{A})$ with compact support and for $f, g \in C_c(G, r^*(\mathcal{A}))$, set

$$f * g(\gamma) := \int f(\eta) \sigma_\eta(g(\eta^{-1}\gamma)) d\lambda^{r(\gamma)}(\eta)$$

and set

$$f^*(\gamma) = \sigma_\gamma(f(\gamma^{-1})^*).$$

Then, with respect to these operations and pointwise addition and scalar multiplication, $C_c(G, r^*(\mathcal{A}))$ becomes a topological $*$ -algebra in the inductive limit topology to which Renault's disintegration theorem [16] applies. The enveloping C^* -algebra of $C_c(G, r^*(\mathcal{A}))$ is called the *crossed product of G acting on $\mathcal{A}$* and is denoted $C^*(G; \mathcal{A})$ [16]. The basic problem we study in this note is

Question 1. Under what circumstances on G and $\mathcal{A}$ is $C^*(G; \mathcal{A})$ a continuous trace C^* -algebra?

Date: 20 May 2002.

¹We follow [4] for the general theory of Banach- and C^* -bundles. However, we adopt the increasingly popular convention that bundles are to be denoted by calligraphic letters. The fibres in a given bundle are then denoted by the corresponding roman letter. Thus if $\mathcal{A}$ is a bundle of C^* -algebras, say, over a space X , then the fibre over $x \in X$ will be denoted $A(x)$.

The first systematic investigation into questions of this type that we know of is Green's pioneering study [5]. There, Green deals with the case where G is the transformation group groupoid obtained by letting a locally compact *group* H act on a locally compact Hausdorff space X and where $\mathcal{A}$ is the trivial line bundle over $G^{(0)} = X$. His principal result asserts that if H acts freely on X , then $C^*(G; \mathcal{A})$ is continuous trace if and only if the action of H is proper. In this event, in fact, $C^*(G; \mathcal{A})$ is strongly Morita equivalent to $C_0(X/H)$.

Another precedent to this investigation is the work of the second two authors in [10]. Here, the hypothesis is that G is a *principal* groupoid and the bundle $\mathcal{A}$ is again the trivial line bundle. It was shown that $C^*(G; \mathcal{A}) = C^*(G)$ has continuous trace if and only if the action of G on $G^{(0)}$ is proper. In this event, again, $C^*(G)$ is strongly Morita equivalent to $C_0(G^{(0)}/G)$.

The first example where the bundle $\mathcal{A}$ is nontrivial was considered by Raeburn and Rosenberg in [14]. They considered a locally compact *group* acting on a continuous trace C^* -algebra A and showed that if the natural action of G on the spectrum of A , $\hat{A}$, is free and proper, then the cross product C^* -algebra $A \rtimes G$ has continuous trace. In [13], Olesen and Raeburn proved a conditioned converse: if the group G is abelian and acts freely on $\hat{A}$, then if $A \rtimes G$ is continuous trace, the action of G on $\hat{A}$ must be proper. Quite recently, Deicke [2] used non-abelian duality theory to remove the hypothesis that G is abelian. Thus, the best result in this direction is: If G acts on A yielding a free action on $\hat{A}$, then $A \rtimes G$ is continuous trace if and only if the action of G on $\hat{A}$ is proper.

Our objective in this note, Theorem 1, is to prove a result that contains all of these examples as special cases — and much more, as well. It is based on two hypotheses and some ancillary considerations that we will elaborate. The first hypothesis is

Hypothesis 1. The C^* -algebra $C_0(G^{(0)}, \mathcal{A})$ is continuous trace.

Here, $C_0(G^{(0)}, \mathcal{A})$ denotes the C^* -algebra of continuous sections of the bundle $\mathcal{A}$ that vanish at infinity on $G^{(0)}$. An hypothesis on the bundle $\mathcal{A}$ of this nature is natural and reasonable, in view of the fact that in the trivial case $G = G^{(0)}$, we have $C^*(G; \mathcal{A}) \cong C_0(G^{(0)}, \mathcal{A})$. Furthermore, we note that a compact group can act on an antiliminal C^* -algebra in such a way that the crossed product is continuous trace [17].

To state our second hypothesis, we need a couple of remarks about the spectrum of $C_0(G^{(0)}, \mathcal{A})$. We shall denote it by X throughout this note. Observe that X may be expressed as the disjoint union of spectra $\coprod_{u \in G^{(0)}} \hat{A}(u)^\wedge$ and the natural projection $\hat{p}$ from X to $G^{(0)}$ is continuous and open [8, 12]. The groupoid G acts on X (using the map $\hat{p}$) as follows. If $x \in X$, we shall write $x = [\pi_x]$ in order to specify a particular irreducible representation in the equivalence class represented by x . Then if $X * G$ denotes the space $\{(x, \gamma) \in X \times G \mid \hat{p}(x) = r(\gamma)\}$ and if $(x, \gamma) \in X * G$, then $x \cdot \gamma$ is defined to be $[\pi_x \circ \sigma_\gamma]$. The fact that the action of G on X is well defined and is continuous is easily checked (as in [15, Lemma 7.1] for example). Our second hypothesis is

Hypothesis 2. The action of G on X is free.

For $x \in X$, we shall write $\mathcal{K}(x)$ for the quotient $\mathcal{A}(\hat{p}(x))/\ker(\pi_x)$. Then $\mathcal{K}(x)$ is well defined (i.e., it is independent of the choice of π_x) and is an elementary C^* -algebra. Hypothesis 1 guarantees that the collection $\{\mathcal{K}(x)\}_{x \in X}$ may be given the

structure of an elementary C^* -algebra bundle over X satisfying Fell's condition [3, Proposition 10.5.8]. As a result, we find that $C_0(G^{(0)}, \mathcal{A})$ is naturally isomorphic to $C_0(X, \mathcal{K})$. The action of G on X induces one on $\mathcal{K}$ that we shall use. We find it preferable to express this in terms of the *action groupoid* $X * G$ defined as follows.

As a set, $X * G := \{(x, \gamma) \mid \hat{p}(x) = r(\gamma)\}$ and the groupoid operations are defined by the formulae

$$(x, \alpha)(x\alpha, \beta) := (x, \alpha\beta) \text{ and} \\ (x, \alpha)^{-1} := (x\alpha, \alpha^{-1}),$$

$(x, \alpha), (x, \beta) \in X * G$. Note, in particular, that the unit space of $X * G$ may be identified with X via: $(x, \hat{p}(x)) \longleftrightarrow x$. Note, too, that the range and source maps on $X * G$, denoted $\tilde{r}$ and $\tilde{s}$, are given by the equations $\tilde{r}(x, \alpha) = (x, r(\alpha))$ and $\tilde{s}(x, \alpha) = (x\alpha, s(\alpha))$. The groupoid $X * G$ in the product topology is clearly locally compact, Hausdorff, and separable. It has a Haar system $\{\tilde{\lambda}^x\}_{x \in X}$ given by the formula $\tilde{\lambda}^x = \delta_x \times \lambda^{\hat{p}(x)}$. Observe that the action of G on X is free (resp. proper) iff $X * G$ is principal (resp. proper).

The groupoid $X * G$ acts on $\mathcal{K}$ via the formula

$$\tilde{\sigma}_{(x, \gamma)}(k) := \sigma_\gamma(a) + \ker(\pi_x),$$

where $k = a + \ker \pi_{x\gamma}$ lies in $K(\tilde{s}(x, \gamma))$. Note that this action is well defined since $\pi_x \circ \sigma_\gamma = \pi_{x\gamma}$. We promote this action of $X * G$ on $\mathcal{K}$ to one on $X * \mathcal{K} := \{(x, k) \mid k \in K(x)\}$: $(x, k) \cdot (x, \gamma) := (x \cdot \gamma, \tilde{\sigma}_{(x, \gamma)}^{-1}(k))$. If the action of G on X is free and proper, then the action of $X * G$ on $X * \mathcal{K}$ is also free and proper. In this case, we write $\mathcal{K}^X$ for the quotient space $X * \mathcal{K} / X * G$. Then $\mathcal{K}^X$ is naturally a bundle of elementary C^* -algebras over X/G . In fact, using the methods of Theorem 1.1 of [14], it is easy to see that $\mathcal{K}^X$ satisfies Fell's condition. Thus, in particular, $C_0(X/G, \mathcal{K}^X)$ has continuous trace, if G acts on X freely and properly.

With these preliminaries at our disposal, we are able to state the main result of this paper as

Theorem 1. *Under Hypotheses 1 and 2, $C^*(G; \mathcal{A})$ has continuous trace if and only if the action of G on X is proper. In this event, $C^*(G; \mathcal{A})$ is strongly Morita equivalent to $C_0(X/G, \mathcal{K}^X)$, where $\mathcal{K}^X$ is the elementary C^* -bundle over X/G , satisfying Fell's condition, that was just defined.*

2. SUFFICIENCY IN THEOREM 1

First we reduce the proof of Theorem 1 to the case when G is a principal groupoid. This reduction is accomplished with the aid of

Theorem 2. *In the notation established above, $C^*(G; \mathcal{A})$ is isomorphic to $C^*(X * G; \mathcal{K})$.*

Proof. For $u \in G^{(0)}$, identify $A(u)$ with $C_0(\hat{p}^{-1}(u), \mathcal{K})$. Also, set $C_{cc}(G, r^*(\mathcal{A})) := \{f \in C_c(G, r^*(\mathcal{A})) \mid (x, \gamma) \rightarrow \|f(\gamma)(x)\| \text{ has compact support in } X * G\}$. Recall that $r^*(\mathcal{A}) = \{(\gamma, a) \mid a \in A(r(\gamma))\}$ and, likewise, $\tilde{r}^*(\mathcal{K}) = \{((x, \gamma), a) \mid a \in K(x)\}$. So, in the definition of $C_{cc}(G, r^*(\mathcal{A}))$, $f(\gamma)$, which nominally is in $A(r(\gamma))$, is to be viewed in $C_0(\hat{p}^{-1}(u), \mathcal{K})$. In particular, $f(\gamma)(x)$ lies in $K(x)$, when $\hat{p}(x) = r(\gamma)$. Now it is straightforward to check that $C_{cc}(G, r^*(\mathcal{A}))$ is a subalgebra of

$C_c(G, r^*(\mathcal{A}))$ that is dense in $C_c(G, r^*(\mathcal{A}))$ in the inductive limit topology. Further, if we define $\Psi : C_c(X * G, \tilde{r}^*(\mathcal{K})) \rightarrow C_{cc}(G, r^*(\mathcal{A}))$ by the formula

$$[\Psi(f)(\gamma)](x) = f(x, \gamma),$$

$f \in C_c(X * G, \tilde{r}^*(\mathcal{K}))$, then Ψ is an algebra $*$ -homomorphism that is continuous in the inductive limit topologies on $C_c(X * G, \tilde{r}^*(\mathcal{K}))$ and $C_{cc}(G, r^*(\mathcal{A}))$. Hence by Renault's disintegration theorem [16, Theorem 4.1], Ψ extends to a C^* -homomorphism from $C^*(X * G; \mathcal{K})$ into $C^*(G; \mathcal{A})$.

The inverse $\Phi : C_{cc}(G, r^*(\mathcal{A})) \rightarrow C_c(X * G, \tilde{r}^*(\mathcal{K}))$ to Ψ is given formally by the formula

$$\Phi(f)(x, \gamma) = f(\gamma)(x),$$

$f \in C_{cc}(G, r^*(\mathcal{A}))$; i.e., at the level of C_c , $\Phi \circ \Psi$ and $\Psi \circ \Phi$ are the identity maps on the appropriate algebras. The problem is that *a priori* Φ is not continuous in the inductive limit topologies. However, Φ is manifestly continuous with respect to the so-called L^I -norms on $C_{cc}(G, r^*(\mathcal{A}))$ and $C_c(X * G, \tilde{r}^*(\mathcal{K}))$, where for $f \in C_{cc}(G, r^*(\mathcal{A}))$, the L^I -norm of f is

$$\max \left\{ \sup_{u \in G^{(0)}} \int \|f(\gamma)\|_{A(u)} d\lambda^u, \sup_{u \in G^{(0)}} \int \|f^*(\gamma)\|_{A(u)} d\lambda^u \right\},$$

and similarly for $C_c(X * G, \tilde{r}^*(\mathcal{K}))$. Since every representation of one of these algebras is continuous in the L^I -norm by Renault's disintegration theorem [16, Theorem 4.1], we conclude that Φ extends to a C^* -homomorphism from $C^*(G; \mathcal{A})$ to $C^*(X * G; \mathcal{K})$ that is the inverse of Ψ . $\square$

On the basis of Theorem 2 we may and shall assume from now on that $X = G^{(0)}$ and that $A(u)$ is an elementary C^ -algebra for every $u \in G^{(0)}$.*

Proof of the Sufficiency. If G acts freely and properly on $G^{(0)}$, then $G^{(0)}$ is an equivalence between G and the quotient space $G^{(0)}/G$ (viewed as a cotrivial groupoid) in the sense of [9]. Further, if $G^{(0)} * \mathcal{A} := \{(u, a) \mid a \in A(u)\}$, then $G^{(0)} * \mathcal{A}$ serves as a bundle of Morita equivalences between $\mathcal{A}$ and $\mathcal{A}^{G^{(0)}}$ in the sense of [7]. (See [16], too.) Indeed, recall that $\mathcal{A}^{G^{(0)}}$ is the quotient $(G^{(0)} * \mathcal{A})/G$, where $(r(\gamma), a) \cdot \gamma = (s(\gamma), \sigma_\gamma^{-1}(a))$. Then the $\mathcal{A}$ -valued inner product on $G^{(0)} * \mathcal{A}$ is given by the formula

$$\langle (u, a), (u, b) \rangle_{\mathcal{A}} = a^*b,$$

while the $\mathcal{A}^{G^{(0)}}$ -valued inner product on $G^{(0)} * \mathcal{A}$ is given by the formula

$${}_{\mathcal{A}^{G^{(0)}}} \langle (u, a), (u, b) \rangle = [u, ab^*],$$

where $[u, ab^*]$ denotes the image of (u, ab^*) in $\mathcal{A}^{G^{(0)}}$. By Corollaire 5.4 of [16],

$$C^*(G; \mathcal{A}) \text{ is Morita equivalent to } C_0(G^{(0)}/G, \mathcal{A}^{G^{(0)}}).$$

This second algebra is a continuous-trace C^* -algebra since $\mathcal{A}^{G^{(0)}}$ is a bundle of elementary C^* -algebras satisfying Fell's condition, as we noted earlier. $\square$

3. NECESSITY IN THEOREM 1

The proof of the necessity in Theorem 1 is modeled on the proofs in [10] and [11], which in turn are inspired by ideas in [5]. There are, however, a number of new difficulties that must be overcome.

For each $u \in G^{(0)}$, we fix an irreducible representation π_u of $A(u)$ on a Hilbert space H_u . Since each $A(u)$ is elementary the π_u 's are unique up to unitary equivalence. We define L^u to be the representation of $C^*(G; \mathcal{A})$ on the Hilbert space $L^2(\lambda_u) \otimes H_u$ according to the formula

$$(1) \quad L^u(f)\xi(\gamma) = \int \pi_u \circ \sigma_\gamma^{-1}(f(\gamma\alpha))\xi(\alpha^{-1}) d\lambda^u(\alpha),$$

where $f \in C_c(G, r^*(\mathcal{A}))$ and $\xi \in L^2(\lambda_u) \otimes H_u$. (Recall that λ_u is the image of λ^u under inversion.) Thus, L^u is the representation of $C^*(G; \mathcal{A})$ induced by the irreducible representation π_u viewed as a representation of $C_0(G^{(0)}, \mathcal{A})$. This implies, in particular, that replacing π_u by a unitarily equivalent representation does not affect the unitary equivalence class of L^u . The following lemma and corollary capture the salient features of the L^u that we shall use.

Lemma 3. *Under the hypothesis that the action of G on $G^{(0)}$ is free (and Hypothesis 1 on $\mathcal{A}$), the following assertions hold:*

- (a) *Each representation L^u is irreducible.*
- (b) *L^u is unitarily equivalent to L^v if and only if u and v lie in the same orbit.*
- (c) *The map $u \rightarrow L^u$ is continuous.*

Proof. The proof follows the lines of the arguments in [10, Lemma 2.4 and Proposition 2.5]. Only minor changes need to be made to accommodate the presence of $\mathcal{A}$. The key point is that L^u is unitarily equivalent to the representation R^u of $C^*(G; \mathcal{A})$ defined by the formula

$$R^u(f)\xi(\gamma \cdot u) = \int \pi_u \circ \sigma_\gamma^{-1}(f(\gamma\alpha))\xi(\alpha^{-1} \cdot u) d\lambda^u(\alpha),$$

$f \in C_c(G, r^*(\mathcal{A}))$, $\xi \in L^2([u], \mu_{[u]}) \otimes H_u$, where $[u]$ denotes the orbit of u and $\mu_{[u]}$ is the image of λ_u under the map $r|s^{-1}(u)$. The fact that the action of G on $G^{(0)}$ is free (i.e., G is a principal groupoid) implies that $r|s^{-1}(u)$ is a bijection between $s^{-1}(u)$ and $[u]$. It is a Borel isomorphism, of course, because of our separability hypotheses and the fact that $r|s^{-1}(u)$ is continuous.

The value of R^u for us lies in the fact that it is evident how to express R^u as the integrated form of a representation of $(G, \mathcal{A})$ in the sense of [16, Definition 3.4]. The measure class on $G^{(0)}$ is, of course, that determined by $\mu_{[u]}$ and the Hilbert bundle $\mathcal{H}$ is the constant bundle determined by H_u over the orbit of u , i.e.,

$$H(v) = \begin{cases} \{v\} \times H_u & v \in [u] \\ 0 & \text{otherwise} \end{cases}.$$

Thus, $\int^\oplus H(v) d\mu_{[u]}$ is identified with $L^2([u], \mu_{[u]}) \otimes H_u$ in the standard fashion. The groupoid G is represented on $\mathcal{H}$ according to the formula

$$U_\gamma((s(\gamma), \xi)) = (r(\gamma), \xi),$$

$\xi \in H_u$, $s(\gamma) \in [u]$, i.e., $\{U_\gamma\}_{\gamma \in G}$ is just the translation representation, and $\mathcal{A}$ is represented on $\mathcal{H}$ according to the formula

$$a \cdot (v, \xi) = (v, \pi_u \circ \sigma_\gamma(a)\xi),$$

$a \in A(v)$, $(v, \xi) \in H(v)$, where γ is the unique element in G with source v and range u .

Observe that the $C_0(G^{(0)}, \mathcal{A})$ acts as multipliers on $C^*(G; \mathcal{A})$ according to the formula $\Phi \cdot f(\gamma) = \Phi(r(\gamma))f(\gamma)$ for $\Phi \in C_0(G^{(0)}, \mathcal{A})$ and $f \in C_c(G, r^*(\mathcal{A}))$. The extension $\tilde{R}^u$ of R^u to the multiplier algebra $C^*(G; \mathcal{A})$ represents $C_0(G^{(0)}, \mathcal{A})$ on $L^2([u], \mu_{[u]}) \otimes H_u$ via the equation

$$\tilde{R}^u(\Phi)\xi(v) = \pi_u \circ \sigma_\gamma(\Phi(v))\xi(v),$$

again, where γ is the unique element in G with source v and range u . It is clear from this that the weak closure of the algebra $\tilde{R}^u(C_0(G^{(0)}, \mathcal{A}))$ is the full algebra of decomposable operators on $L^2([u], \mu_{[u]}) \otimes H_u$. It follows that any projection that commutes with $R^u(C^*(G; \mathcal{A}))$ must be diagonal. On the other hand, it follows from the definition of the representation of G , $\{U_\gamma\}_{\gamma \in G}$, that a diagonal operator commuting with $R^u(C^*(G; \mathcal{A}))$ must commute with $\{U_\gamma\}_{\gamma \in G}$, and therefore must be constant a.e. $\mu_{[u]}$. This proves that R^u , and hence L^u , is irreducible.

If u and v lie in the same orbit, it is clear that translation by the (unique) γ with source v and range u implements an equivalence between L^u and L^v . On the other hand, if u and v lie in different orbits, then L^u and L^v are disjoint. Indeed, the representations N_u and N_v of $C_0(G^{(0)})$ obtained by restricting $\tilde{R}^u$ and $\tilde{R}^v$ to $C_0(G^{(0)})$, viewed as a subalgebra of $M(C^*(G; \mathcal{A}))$, are supported on the disjoint sets $[u]$ and $[v]$. Arguing just as we did in the proof of [10, Proposition 2.5], using [18, Lemma 4.15], we conclude L^u and L^v are disjoint.

Finally, to see that the map $u \mapsto L^u$ is continuous, observe that Hypothesis 1 guarantees that for each point $u \in G^{(0)}$ we can find a neighborhood $\mathcal{V}_u$ of u on which the H_v 's can be chosen to be the fibres of a (topological) Hilbert bundle $\tilde{\mathcal{H}}$ and on which we can choose the π_v 's so that for any section $\Phi \in C_0(G^{(0)}, \mathcal{A})$ that is supported on $\mathcal{V}_u$ and any two C_0 -sections of $\tilde{\mathcal{H}}$ over $\mathcal{V}_u$, ξ and η , the function $v \mapsto (\pi_v(\Phi(v))\xi(v) \mid \eta(v))_{H_v}$ is continuous. It follows from the continuity of the Haar system that given such sections ξ and η of $\tilde{\mathcal{H}}$ and any two functions g and h in $C_c(G)$, the function $v \mapsto (L^v(f)(g \otimes \xi) \mid (h \otimes \eta))$ (where the inner products are taken in $L^2(\lambda_v) \otimes H_v$) is continuous for all $f \in C_c(G; r^*(\mathcal{A}))$. This shows that the map $u \mapsto L^u$ is continuous. $\square$

Corollary 4. *Assume that G is principal and that $\mathcal{A}$ is an elementary C^* -bundle over $G^{(0)}$, satisfying Fell's condition, on which G acts. If $C^*(G; \mathcal{A})$ has continuous trace, then the map that sends $u \in G^{(0)}$ to the unitary equivalence class of L^u defines a continuous open surjection of $G^{(0)}$ onto $C^*(G; \mathcal{A})^\wedge$ that is constant on G -orbits. In particular, orbits are closed and $G^{(0)}/G$ is homeomorphic to $C^*(G; \mathcal{A})^\wedge$.*

Proof. The proof is also essentially the same as the proof in [10, Proposition 2.5]. Here is an outline. Write Ψ for the map $u \mapsto [L^u]$. Then by Lemma 3, Ψ is continuous and constant on G -orbits. Thus Ψ passes to continuous map on $G^{(0)}/G$ with the quotient topology (no matter how bad that might be). Since, however, $C^*(G; \mathcal{A})^\wedge$ is Hausdorff by hypothesis, we conclude that $G^{(0)}/G$ is Hausdorff.

Suppose that L is an irreducible representation of $C^*(G; \mathcal{A})$ and let M be the representation of $C_0(G^{(0)})$ obtained by extending L to the multiplier algebra of $C^*(G; \mathcal{A})$ and then restricting to $C_0(G^{(0)})$. The kernel J of M is the set of functions in $C_0(G^{(0)})$ that vanish on a closed set F in $G^{(0)}$. Then F is easily seen to be invariant. Indeed, one may do this directly or use the fact that it supports the

quasi-invariant measure associated to the disintegrated form of L guaranteed by [16]. Further, since L is irreducible, F cannot be expressed as the union of two disjoint, closed, G -invariant sets. Since the quotient map from $G^{(0)}$ to $G^{(0)}/G$ is continuous and open, we may apply the lemma on page 222 of [6] to conclude that F is an orbit closure. Since we now know that orbits are closed, F is, in fact, an orbit. Thus L factors through $C^*(G|_{[u]}; \mathcal{A})$. However, $G|_{[u]}$ is a *transitive* principal groupoid. A little reflection, using Theorem 3.1 of [9], reveals that every irreducible representation of $C^*(G|_{[u]}; \mathcal{A})$ is unitarily equivalent to L^u . $\square$

We now assume the action of G on $G^{(0)}$ is not proper and use Lemma 2.6 of [10] to choose a sequence $\{\gamma_n\} \subseteq G$ such that $\gamma_n \rightarrow \infty$ in the sense that $\{\gamma_n\}$ eventually escapes each compact subset of G , and such that $r(\gamma_n), s(\gamma_n) \rightarrow z$, for some $z \in G^{(0)}$. We shall fix this sequence for the remainder of the proof. We also choose a relatively compact neighborhood U of $z \in G^{(0)}$ and a section g in the Pedersen ideal of $C_0(G^{(0)}, \mathcal{A})$ such that g is non-negative, compactly supported, and satisfies $\text{tr}(\pi_u(g(u))) \equiv 1$ on U . The fact that $\mathcal{A}$ satisfies Fell's condition guarantees that such choices are possible.

With these ingredients fixed, we want to build a special neighborhood E of z in G , following the analysis on pages 236–238 of [10]. First observe, as we have above, that since G is principal, r maps G_z bijectively onto $[z]$ while s maps G^z bijectively onto $[z]$, where, recall, $[z]$ denotes the orbit of z . Since $[z]$ is closed by Corollary 4 while r is continuous and open on G , we see that r maps G_z homeomorphically onto $[z]$. Likewise, s maps G^z homeomorphically onto $[z]$. Also, since G is principal, multiplication induces a homeomorphism between $G_z \times G^z$ and $G|_{[z]}$. Let N be the closed support of g , a compact subset of $G^{(0)}$, and set $F_z := G_z \cap r^{-1}([z] \cap N)$ and $F^z := G^z \cap s^{-1}([z] \cap N)$, obtaining compact subsets of G_z and G^z , respectively. Then we see that if $\gamma \in G|_{[z]}$ and if $g(s(\gamma)) \neq 0$ and $g(r(\gamma)) \neq 0$, then $\gamma \in F_z F^z$.

According to Lemma 2.7 of [10], we may select symmetric, conditionally compact open neighborhoods W_0 and W_1 of $G^{(0)}$ such that $\overline{W_0} \subseteq W_1$. (Recall that a neighborhood W of $G^{(0)}$ is conditionally compact in case VW and WV are relatively compact subsets of G for each relatively compact subset V in G .) We select such a pair, as we may, with the additional property that $F_z F^z \subseteq W_0 z W_0$. Then from the preceding paragraph, we see that if $\gamma \notin W_0 z W_0$, then either $g(s(\gamma)) = 0$ or $g(r(\gamma)) = 0$.

By construction,

$$\overline{W_1}^7 z \setminus W_0 z \subseteq r^{-1}(G^{(0)} \setminus N).$$

So we may find relatively compact open neighborhoods V_0 and V_1 of z in G so that $V_0 \subseteq W_0$, $\overline{V_0} \subseteq V_1$, and so that

$$\overline{W_1}^7 \overline{V_1} \setminus W_0 V_0 \subseteq r^{-1}(G^{(0)} \setminus N).$$

With these V_0 and V_1 so chosen, the special open neighborhood E of z in G that we want is defined to be $E := W_0 V_0 W_0$.

Observe that we have

$$\overline{W_1}^7 \overline{V_1} \overline{W_1}^7 \setminus E = \overline{W_1}^7 \overline{V_1} \overline{W_1}^7 \setminus W_0 V_0 W_0 \subseteq r^{-1}(G^{(0)} \setminus N).$$

Set

$$g^1(\gamma) := \begin{cases} g(r(\gamma)), & \gamma \in \overline{W_1}^7 \overline{V_1} \overline{W_1}^7 \\ 0, & \gamma \notin E \end{cases}.$$

Then, since $\overline{W_1^7 V_1} \overline{W_1^7} \setminus E \subseteq r^{-1}(G^{(0)} \setminus N)$, g^1 is a continuous section of $r^*(\mathcal{A})$ on G that vanishes outside E .

Observe the following containment relations among relatively compact sets: $E^2 = W_0 V_0 W_0^2 V_0 W_0 \subseteq W_0^4 V_0 W_0^4 \subseteq \overline{W_0^4 V_0 W_0^4} \subseteq W_1^4 V_1 W_1^4$. Hence, we may find a compactly supported function b on G such that $0 \leq b(\gamma) \leq 1$ for all γ , $b \equiv 1$ on E^2 and $b \equiv 0$ off $W_1^4 V_1 W_1^4$. Replacing b by $\frac{b+b^*}{2}$, if necessary, we may assume that b is a selfadjoint element of the convolution algebra of scalar-valued functions $C_c(G)$.

Define $F(\gamma) := g(r(\gamma))\sigma_\gamma(g(s(\gamma)))b(\gamma)$. By our choices of g and b , F belongs to $C_c(G, r^*(\mathcal{A}))$, F is selfadjoint and

$$L^u(F)\xi(\gamma) = \pi_u(\sigma_\gamma^{-1}(g(r(\gamma)))) \int \pi_u(\sigma_\alpha^{-1}(g(r(\alpha))))b(\gamma\alpha^{-1})\xi(\alpha) d\lambda_u(\alpha)$$

for all u by the definition of L^u (cf. (1)). Let $P_{u,1}$ be the projection onto $\mathcal{E}_{u,1} := L^2(G_u \cap E) \otimes H_u$ and let $P_{u,2}$ be the projection onto the orthocomplement, $\mathcal{E}_{u,2} := L^2(G_u \setminus E) \otimes H_u$. Then, if $P_{u,1}\xi = \xi$, we see that

$$\begin{aligned} L^u(F)\xi(\gamma) &= \pi_u(\sigma_\gamma^{-1}(g(r(\gamma)))) \int_{G_u \cap E} \pi_u(\sigma_\alpha^{-1}(g(r(\alpha))))\xi(\alpha) d\lambda_u(\alpha) \\ &= \pi_u \circ \sigma_\gamma^{-1}(g^{(1)}(\gamma)) \int_{G_u \cap E} \pi_u \circ \sigma_\alpha^{-1}(g^{(1)}(\alpha))\xi(\alpha) d\lambda_u(\alpha) \end{aligned}$$

for all $\gamma \in G_u \cap E$ because b is identically 1 on E^2 . However, by definitions of E and g^1 , the equation persists when $\gamma \in G_u \setminus E$, yielding 0. Thus, $P_{u,1}$ commutes with $L^u(F)$. Moreover, when $u = z$, these formulas show that $L^z(F)P_{z,1} = L^z(F)$.

We now want to show that $L^u(F)P_{u,1} \geq 0$ and we want to analyze the trace, $\text{tr}(L^u(F)P_{u,1})$. However, when ξ is in the range of $P_{u,1}$, the formula for $L^u(F)\xi$ shows that

$$\begin{aligned} (L^u(F)\xi \mid \xi) &= \iint (\pi_u \circ \sigma_\gamma^{-1}(g^{(1)}(\gamma))\pi_u \circ \sigma_\alpha^{-1}(g^{(1)}(\alpha))\xi(\alpha), \xi(\gamma)) d\lambda_u(\alpha) d\lambda_u(\gamma) \\ &= \iint (\pi_u \circ \sigma_\alpha^{-1}(g^{(1)}(\alpha))\xi(\alpha), \pi_u \circ \sigma_\gamma^{-1}(g^{(1)}(\gamma))\xi(\gamma)) d\lambda_u(\alpha) d\lambda_u(\gamma) \\ &\geq 0. \end{aligned}$$

As for the trace, observe that if K_u is defined by the formula

$$K_u(\gamma, \eta) = \pi_u(\sigma_\gamma^{-1}(g^1(\gamma))\sigma_\eta^{-1}(g^1(\eta)))$$

on $G_u \times G_u$, then our calculations show that K_u is continuous, positive semi-definite and supported on $(G_u \cap E) \times (G_u \cap E)$, and that $(L^u(F)P_{u,1}\xi \mid \zeta) = \iint (K_u(\gamma, \eta)\xi(\eta), \zeta(\gamma)) d\lambda_u(\eta) d\lambda_u(\gamma)$. Consequently, we may use Duflo's generalization of Mercer's theorem, [1, Proposition 3.1.1], and the fact that $K_u(\gamma, \gamma) = \pi_u \circ \sigma_\gamma^{-1}(g(r(\gamma)))^2$ to conclude that $\text{tr}(L^u(F)P_{u,1}) =$

$$\int_{G_u \cap E} \text{tr}(\pi_u \circ \sigma_\gamma^{-1}(g(r(\gamma)))^2) d\lambda_u(\gamma).$$

By our choice of g , this expression is continuous in u and when $u = z$ yields the value $\text{tr}(L^z(F))$.

We will show that there is a positive number a such that

$$(2) \quad \|(L^{s(\gamma_n)}(F)P_{s(\gamma_n),2})^+\| \geq 2a$$

eventually, where $(L^{s(\gamma_n)}(F)P_{s(\gamma_n),2})^+$ denotes the positive part of the selfadjoint operator $L^{s(\gamma_n)}(F)P_{s(\gamma_n),2}$. Therefore, eventually

$$(\text{the largest eigenvalue of } L^{s(\gamma_n)}(F)P_{s(\gamma_n),2}) \geq 2a.$$

Assume that we have shown this, and set

$$q(t) = \begin{cases} 0, & t \leq a \\ 2(t-a), & a \leq t \leq 2a \\ t, & 2a \end{cases}$$

Then $q(F)$ is a positive element in the Pedersen ideal of $C^*(G, \mathcal{A})$, and so the function $u \rightarrow \text{tr}(L^u(q(F))) =$

$$\text{tr}(L^u(q(F))P_{u,1}) + \text{tr}(L^u(q(F))P_{u,2})$$

is finite and continuous in u , with value $\text{tr}(L^z(q(F)))$ at $u = z$. (Recall that $L^z(F) = L^z(F)P_{z,1}$ and so $L^z(q(F)) = L^z(q(F))P_{z,1}$.) On the other hand, we showed that $L^u(F)P_{u,1}$ is positive. Since $P_{u,1}$ commutes with $L^u(F)$, $L^u(F)P_{u,1} = L^u(F^+)P_{u,1}$. But also we showed that $u \rightarrow \text{tr}(L^u(F)P_{u,1})$ is continuous at z . Consequently, so is $u \rightarrow \text{tr}(L^u(F^+)P_{u,1})$. Since $q(F) \leq F^+$, the function $u \rightarrow \text{tr}(L^u(q(F))P_{u,1})$ is continuous by Lemma 4.4.2(i) in [3], with value $\text{tr}(L^z(q(F))) = \text{tr}(L^z(q(F))P_{z,1})$ at $u = z$, also. Therefore

$$\lim_{u \rightarrow z} \text{tr}(L^u(q(F))P_{u,2}) = 0.$$

Since the largest eigenvalue of $L^{s(\gamma_n)}(F)P_{s(\gamma_n),2} \geq 2a$, the largest eigenvalue of $L^{s(\gamma_n)}(q(F))P_{s(\gamma_n),2} \geq 2a$ also. Consequently,

$$\liminf_n \text{tr}(L^{s(\gamma_n)}(q(F))P_{s(\gamma_n),2}) \geq 2a.$$

This contradiction will complete the proof.

So we will finish by verifying the asserted inequality (2). To this end, choose an open neighborhood of z in G , V_2 , that is contained in V_0 and choose a conditionally compact neighborhood Y of $G^{(0)}$ such that if $v \in V_2$, then r maps Yv into U . Without loss of generality, we may assume that $Y \subseteq W_0$. Observe that if $\gamma_n \notin \overline{W_1}^2 \overline{V_1} \overline{W_1}^2$ then for $\gamma \in Y\gamma_n$, $\gamma \notin E$. Indeed, if $\gamma = \gamma'\gamma_n \in E \cap Y\gamma_n$, then $\gamma_n \in (\gamma')^{-1}E \subseteq W_0^2 V_0 W_0 \subseteq \overline{W_1}^2 \overline{V_1} \overline{W_1}^2$ contrary to assumption. So, since $r(\gamma_n)$ and $s(\gamma_n)$ are tending to z , while γ_n eventually escapes $\overline{W_1}^2 \overline{V_1} \overline{W_1}^2$, we can conclude that for n sufficiently large, whenever γ lies in $Y\gamma_n$, then $\gamma \notin E$ while $r(\gamma)$ and $s(\gamma)$ lie in U . From now on, we will assume that n is sufficiently large so that these conditions are satisfied.

Next observe that since for each n , the map $\gamma \rightarrow \pi_{s(\gamma_n)} \circ \sigma_\gamma^{-1}(g(r(\gamma)))$ defines a continuous family of rank 1 projections on the Hilbert space $H_{s(\gamma_n)}$, we can find a Borel family of unit vectors $\gamma \rightarrow v_\gamma^n$ such that $\pi_{s(\gamma_n)} \circ \sigma_\gamma^{-1}(g(r(\gamma)))$ is the rank 1 projection determined by v_γ^n .

Let $h_n(\gamma) = 1_{Y\gamma_n}(\gamma) \times v_\gamma^n$ where $1_{Y\gamma_n}$ denotes the characteristic function of $Y\gamma_n$. Observe that if $\gamma, \alpha \in Y\gamma_n$, then $\gamma\alpha^{-1} \in Y\gamma_n\gamma_n^{-1}Y \subseteq YV_0Y \subseteq W_0V_0W_0 = E$.

Consequently, $b(\gamma\alpha^{-1}) = 1$ and we may calculate to find that if $\gamma \in Y\gamma_n$, then

$$\begin{aligned} L^{s(\gamma_n)}(F)h_n(\gamma) &= \pi_{s(\gamma_n)}(\sigma_\gamma^{-1}(g(r(\gamma)))) \int \pi_{s(\gamma_n)}(\sigma_\alpha^{-1}(g(r(\alpha))))b(\gamma\alpha^{-1})h_n(\alpha) d\lambda_{s(\gamma_n)}(\alpha) \\ &= v_\gamma^n \lambda_{r(\gamma_n)}(Y). \end{aligned}$$

Hence $(L^{s(\gamma_n)}(F)h_n \mid h_n) = \lambda_{r(\gamma_n)}(Y)^2$. However, by our assumption on γ_n , h_n lies in $\mathcal{E}_{s(\gamma_n),2}$, so $((L^{s(\gamma_n)}(F)P_{s(\gamma_n),2})^+h_n \mid h_n) \geq (L^{s(\gamma_n)}(F)P_{s(\gamma_n),2}h_n \mid h_n) = (L^{s(\gamma_n)}(F)h_n \mid h_n) = \lambda_{r(\gamma_n)}(Y)^2$. This shows that $\|(L^{s(\gamma_n)}(F)P_{s(\gamma_n),2})^+\| \geq \lambda_{r(\gamma_n)}(Y)$ provided n is sufficiently large. But the continuity of the Haar system implies that $\liminf \lambda_{r(\gamma_n)}(Y) > 0$, as $n \rightarrow \infty$. This verifies equation (2) and completes the proof of Theorem 1.

REFERENCES

- [1] P. Bernat, N. Conze, M. Duflo, M. Lévy-Nahas, M. Raïs, P. Renouard, and M. Vergne, *Représentations des groupes de Lie résolubles*, Dunod, Paris, 1972, Monographies de la Société Mathématique de France, No. 4.
- [2] Klaus Deicke, *Pointwise unitary coactions on C^* -algebras with continuous trace*, J. Operator Theory **43** (2000), no. 2, 295–327.
- [3] Jacques Dixmier, *C^* -algebras*, North-Holland Mathematical Library, vol. 15, North-Holland, New York, 1977.
- [4] J. M. G. Fell and Robert Doran, *Representations of $*$ -algebras, locally compact groups, and Banach $*$ -algebraic bundles*, vol. I & II, Academic Press, New York, 1988.
- [5] Philip Green, *C^* -algebras of transformation groups with smooth orbit space*, Pacific J. Math. **72** (1977), 71–97.
- [6] ———, *The local structure of twisted covariance algebras*, Acta Math. **140** (1978), 191–250.
- [7] Alexander Kumjian, Paul S. Muhly, Jean N. Renault, and Dana P. Williams, *The Brauer group of a locally compact groupoid*, Amer. J. Math. **120** (1998), 901–954.
- [8] Ru Ying Lee, *On the C^* -algebras of operator fields*, Indiana Univ. Math. J. **25** (1976), 303–314.
- [9] Paul S. Muhly, Jean Renault, and Dana P. Williams, *Equivalence and isomorphism for groupoid C^* -algebras*, J. Operator Theory **17** (1987), 3–22.
- [10] Paul S. Muhly and Dana P. Williams, *Continuous trace groupoid C^* -algebras*, Math. Scand. **66** (1990), 231–241.
- [11] ———, *Continuous trace groupoid C^* -algebras. II*, Math. Scand. **70** (1992), 127–145.
- [12] May Nilsen, *C^* -bundles and $C_0(X)$ -algebras*, Indiana Univ. Math. J. **45** (1996), 463–477.
- [13] Dorte Olesen and Iain Raeburn, *Pointwise unitary automorphism groups*, J. Funct. Anal. **93** (1990), 278–309.
- [14] Iain Raeburn and Jonathan Rosenberg, *Crossed products of continuous-trace C^* -algebras by smooth actions*, Trans. Amer. Math. Soc. **305** (1988), 1–45.
- [15] Iain Raeburn and Dana P. Williams, *Morita equivalence and continuous-trace C^* -algebras*, Mathematical Surveys and Monographs, vol. 60, American Mathematical Society, Providence, RI, 1998.
- [16] Jean Renault, *Représentations des produits croisés d’algèbres de groupoïdes*, J. Operator Theory **18** (1987), 67–97.
- [17] Masamichi Takesaki, *A liminal crossed product of a uniformly hyperfinite C^* -algebra by a compact Abelian automorphism group*, J. Functional Analysis **7** (1971), 140–146.
- [18] Dana P. Williams, *The topology on the primitive ideal space of transformation group C^* -algebras and CCR transformation group C^* -algebras*, Trans. Amer. Math. Soc. **266** (1981), 335–359.

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85287-1804
E-mail address: `ifulman@math.la.asu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IA 52242
E-mail address: `muhly@math.uiowa.edu`

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755-3551
E-mail address: `dana.williams@dartmouth.edu`